arXiv:math/0601609v1 [math.FA] 25 Jan 2006

On the growth of powers of operators with spectrum contained in Cantor sets

AGRAFEUIL Cyril

Abstract

For $\xi \in \left(0, \frac{1}{2}\right)$, we denote by E_ξ the perfect symmetric set associated to ξ, that is

$$E_\xi = \Big\{ \exp\big(2i\pi(1-\xi)\sum_{n=1}^{+\infty} \epsilon_n \xi^{n-1}\big) : \epsilon_n = 0 \text{ or } 1 \quad (n \geq 1) \Big\}.$$

Let s be a nonnegative real number, and T be an invertible bounded operator on a Banach space with spectrum included in E_ξ. We show that if

$$\big\|T^n\big\| = O\big(n^s\big),\ n \to +\infty$$

$$\text{and} \quad \big\|T^{-n}\big\| = O\big(e^{n^\beta}\big),\ n \to +\infty \text{ for some } \beta < \frac{\log\frac{1}{\xi} - \log 2}{2\log\frac{1}{\xi} - \log 2},$$

then for every $\varepsilon > 0$, T satisfies the stronger property

$$\big\|T^{-n}\big\| = O\big(n^{s+\frac{1}{2}+\varepsilon}\big),\ n \to +\infty.$$

This result is a particular case of a more general result concerning operators with spectrum satisfying some geometrical conditions.

1 Introduction

We denote by $\mathbb{T}$ the unit circle and by $\mathbb{D}$ the open unit disk. We shall say that a closed subset E of $\mathbb{T}$ is a K-set if there exists a positive constant c such that for any arc L of $\mathbb{T}$,

$$\sup_{z \in L} d(z, E) \geq c|L|, \tag{K}$$

where $|L|$ denotes the length of the arc L and $d(z, E)$ the distance between z and E. Let E be a K-set. We set

$$\delta(E) = \sup\Big\{\delta \geq 0 : \int_0^{2\pi} \frac{1}{d(e^{it}, E)^\delta} \mathrm{d}t < +\infty\Big\}.$$

We have $\delta(E) \geq \dfrac{\log\frac{1}{1-c}}{\log\frac{2}{1-c}}$ (see [2] section 5, proof of lemma 2 and corollary). E. M. Dyn'kin showed in [2] that condition (K) characterizes the interpolating sets for $\Lambda_s^+(\mathbb{T})$, $s > 0$ (see section 2 for the definition of $\Lambda_s^+(\mathbb{T})$). Let s be a nonnegative real number, and let T be an invertible operator on a Banach space. We show (theorem 2.3) that if the spectrum of T is included in E and if T satisfies

$$\begin{aligned} &\|T^n\| = O\big(n^s\big),\ n \to +\infty \\ \text{and} \quad &\|T^{-n}\| = O\big(e^{n^\beta}\big),\ n \to +\infty \text{ for some } \beta < \frac{\delta(E)}{1+\delta(E)}, \end{aligned}$$

then for every $\varepsilon > 0$, T also satisfies the stronger property

$$\|T^{-n}\| = O\big(n^{s+\frac{1}{2}+\varepsilon}\big),\ n \to +\infty. \tag{1}$$

For $\xi \in \Big(0, \dfrac{1}{2}\Big)$, we denote by E_ξ the perfect symmetric set associated to ξ, that is

$$E_\xi = \Big\{ \exp\big(2i\pi(1-\xi)\sum_{n=1}^{+\infty}\epsilon_n \xi^{n-1}\big) : \epsilon_n = 0 \text{ or } 1 \quad (n \geq 1)\Big\}.$$

We set $b(\xi) = \dfrac{\log\frac{1}{\xi} - \log 2}{2\log\frac{1}{\xi} - \log 2}$. We obtain (as a consequence of theorem 2.3) that if the spectrum of T is included in E_ξ, $\|T^n\| = O\big(n^s\big)$, $n \to +\infty$ and $\|T^{-n}\| = O\big(e^{n^\beta}\big)$, $n \to +\infty$ for some $\beta < b(\xi)$, then T satisfies (1). Notice that J. Esterle showed in [5] that if T is a contraction on a Banach space (respectively on a Hilbert space) with spectrum included in $E_{\frac{1}{q}}$ (respectively included in E_ξ) such that $\|T^{-n}\| = O\big(e^{n^\beta}\big)$, $n \to +\infty$ for some $\beta < b\big(\frac{1}{q}\big)$ (respectively $\beta < b(\xi)$), then $\sup\limits_{n\geq 0}\|T^{-n}\| < +\infty$ (respectively T is an isometry). Here q is an integer greater than or equal to 3.

2 Growth of powers of operators

Let p be a non-negative integer. We denote by $\mathcal{C}^p(\mathbb{T})$ the space of p times continuously differentiable functions on $\mathbb{T}$. We set

$$\mathcal{A}^p(\mathbb{D}) = \Big\{ f \in \mathcal{C}^p(\mathbb{T}) : \widehat{f}(n) = 0 \quad (n < 0)\Big\},$$

$\mathcal{C}^\infty(\mathbb{T}) = \bigcap_{p\geq 0}\mathcal{C}^p(\mathbb{T})$ and $\mathcal{A}^\infty(\mathbb{D}) = \bigcap_{p\geq 0}\mathcal{A}^p(\mathbb{D})$.
Let s be a nonnegative real number, we denote by $[s]$ the nonnegative integer such that $[s] \leq s < [s]+1$. We define the Banach algebra

$$\Lambda_s(\mathbb{T}) = \Big\{ f \in \mathcal{C}^{[s]}(\mathbb{T}) : \sup_{z,z'\in\mathbb{T}} \frac{\big|f^{([s])}(z) - f^{([s])}(z')\big|}{|z-z'|^{s-[s]}} < +\infty \Big\},$$

equiped with the norm $\|f\|_{\Lambda_s} = \|f\|_{\mathcal{C}^{[s]}(\mathbb{T})} + \sup\limits_{z,z'\in\mathbb{T}} \dfrac{\left|f^{([s])}(z) - f^{([s])}(z')\right|}{|z-z'|^{s-[s]}}$. We also define the subalgebra

$$\lambda_s(\mathbb{T}) = \Big\{ f \in \mathcal{C}^{[s]}(\mathbb{T}) : \left|f^{([s])}(z) - f^{([s])}(z')\right| = o\big(|z-z'|^{s-[s]}\big),\ |z-z'| \to 0 \Big\},$$

which we equip with the same norm. We also set

$$\begin{aligned} \Lambda_s^+(\mathbb{T}) &= \Big\{ f \in \Lambda_s(\mathbb{T}) : \widehat{f}(n) = 0 \quad (n<0) \Big\} \\ \text{and } \lambda_s^+(\mathbb{T}) &= \Big\{ f \in \lambda_s(\mathbb{T}) : \widehat{f}(n) = 0 \quad (n<0) \Big\}. \end{aligned}$$

We remark that if s is an integer, $\Lambda_s(\mathbb{T}) = \lambda_s(\mathbb{T}) = \mathcal{C}^s(\mathbb{T})$ and so $\Lambda_s^+(\mathbb{T}) = \lambda_s^+(\mathbb{T}) = \mathcal{A}^s(\mathbb{D})$. We define

$$N_s(E) = \big\{ f \in \Lambda_s(\mathbb{T}) : f_{|E} = \ldots = f_{|E}^{([s])} = 0 \big\},$$

and set $N_s^+(E) = N_s(E) \cap \Lambda_s^+(\mathbb{T})$.

Lemma 2.1. *Let s be a nonnegative real number. Then for all $\varepsilon > 0$, we have the following continuous embedding*

$$\Lambda_{s+\frac{1}{2}+\varepsilon}(\mathbb{T}) \hookrightarrow A_s(\mathbb{T}).$$

Proof. For $s = 0$, this is a result of Bernstein (see [9], p.13). The general case is obtained by the same arguments. Let $\varepsilon > 0$, and set $\tilde{s} = s + \frac{1}{2} + \varepsilon$. Let $f \in \Lambda_{\tilde{s}}(\mathbb{T})$. For $h > 0$, define

$$P(h) = \int_0^{2\pi} \left| f^{([\tilde{s}])}(e^{i(t-h)}) - f^{([\tilde{s}])}(e^{i(t+h)}) \right|^2 \mathrm{d}t.$$

It follows from Parseval equality that

$$P(h) = 8\pi \sum_{n=-\infty}^{+\infty} \left| \widehat{f^{([\tilde{s}])}}(n) \right|^2 \sin^2(nh). \tag{2}$$

Let j_0 be the smallest integer such that $[\tilde{s}] < 2^{j_0}$ and let $j \geq j_0$. It follows from the relation $\widehat{f^{([\tilde{s}])}}(n) = \Big(\prod\limits_{k=1}^{[\tilde{s}]} (n+k) \Big) \widehat{f}(n+[\tilde{s}])$ $(n \in \mathbb{Z})$ and from (2) that there exists a constant $C_1 > 0$ independent of f such that

$$P(h) \quad \geq \quad \frac{4}{C_1^2} \sum_{|n|=2^j}^{2^{j+1}-1} \left| \widehat{f}(n+[\tilde{s}]) \right|^2 (1+|n|)^{2[\tilde{s}]} \sin^2(nh). \tag{3}$$

Using the Cauchy-Schwartz inequality, we have

$$\sum_{|n|=2^j}^{2^{j+1}-1} \left| \widehat{f}(n+[\tilde{s}]) \right| (1+|n|)^s \leq \Big(\sum_{|n|=2^j}^{2^{j+1}-1} \left| \widehat{f}(n+[\tilde{s}]) \right|^2 (1+|n|)^{2[\tilde{s}]} \Big)^{\frac{1}{2}} \Big(\sum_{n=2^j}^{2^{j+1}-1} (1+|n|)^{2s-2[\tilde{s}]} \Big)^{\frac{1}{2}} \tag{4}$$

Set $h = \dfrac{\pi}{3.2^j}$. For all integers n such that $2^j \leq |n| \leq 2^{j+1} - 1$, we have $\dfrac{\pi}{3} \leq |nh| \leq \dfrac{2\pi}{3}$, and so $\sin^2(nh) \geq \dfrac{1}{4}$. So, we deduce from (3) that

$$\Big(\sum_{|n|=2^j}^{2^{j+1}-1} \big|\widehat{f}(n + [\tilde{s}])\big|^2 (1 + |n|)^{2[\tilde{s}]} \Big)^{\frac{1}{2}} \leq C_1 P\big(\frac{\pi}{3.2^j}\big)^{\frac{1}{2}}.$$

Then, as $f \in \Lambda_{\tilde{s}}(\mathbb{T})$, we have

$$P\big(\frac{\pi}{3.2^j}\big)^{\frac{1}{2}} \leq (2\pi)^{\frac{1}{2}} \big\|f\big\|_{\Lambda_{\tilde{s}}} \big(\frac{2\pi}{3.2^j}\big)^{\tilde{s}-[\tilde{s}]},$$

so that

$$\Big(\sum_{|n|=2^j}^{2^{j+1}-1} \big|\widehat{f}(n + [\tilde{s}])\big|^2 (1 + |n|)^{2[\tilde{s}]} \Big)^{\frac{1}{2}} \leq C_1 (2\pi)^{\frac{1}{2}} \big\|f\big\|_{\Lambda_{\tilde{s}}} \big(\frac{2\pi}{3.2^j}\big)^{\tilde{s}-[\tilde{s}]}. \tag{5}$$

Furthermore, there exists a constant $C_2 > 0$ such that

$$\Big(\sum_{|n|=2^j}^{2^{j+1}-1} (1 + |n|)^{2s-2[\tilde{s}]} \Big)^{\frac{1}{2}} \leq C_2 2^{j\left(s-[\tilde{s}]+\frac{1}{2}\right)}. \tag{6}$$

Finally we deduce from (4) and the inequalities (5) and (6) that there exists a constant $C_3 > 0$ independent of f such that for all $j \geq j_0$,

$$\sum_{|n|=2^j}^{2^{j+1}-1} \big|\widehat{f}(n + [\tilde{s}])\big| (1 + |n|)^s \leq 2^{j(s-\tilde{s}+\frac{1}{2})} C_3 \big\|f\big\|_{\Lambda_{\tilde{s}}} = 2^{-\varepsilon j} C_3 \big\|f\big\|_{\Lambda_{\tilde{s}}}.$$

Summing over $j \geq j_0$ these inequalities, we get

$$\sum_{|n|\geq 2^{j_0}} \big|\widehat{f}(n + [\tilde{s}])\big| (1 + |n|)^s \leq \frac{C_3}{1 - 2^{-\varepsilon}} \big\|f\big\|_{\Lambda_{\tilde{s}}},$$

On the other hand, we have $\big|\widehat{f}(n)\big| \leq \big\|f\big\|_{\Lambda_{\tilde{s}}}$ for every $n \in \mathbb{Z}$. So, since j_0 is independent of f, there exists a constant $K > 0$ (independent of f) such that

$$\big\|f\big\|_s \leq K \big\|f\big\|_{\Lambda_{\tilde{s}}}$$

$\square$

Before giving the main theorem of the paper, we need the following lemma.

Lemma 2.2. *Let E be a closed subset of $\mathbb{T}$. We assume that there exists $\delta > 0$ for which $\displaystyle\int_0^{2\pi} \frac{1}{d(e^{it},E)^{\delta}}\mathrm{d}t < +\infty$. Let $\beta < \dfrac{\delta}{1+\delta}$ and let T be an invertible operator on a Banach space with spectrum included in E that satisfies*

$$\begin{aligned} &\big\|T^n\big\| = O\big(n^s\big),\ n \to +\infty \quad (\textit{ for some nonnegative real } s)\\ \textit{and}\quad &\big\|T^{-n}\big\| = O\big(e^{n^{\beta}}\big),\ n \to +\infty, \end{aligned}$$

Then there exists an outer function $f \in \mathcal{A}^{\infty}(\mathbb{D})$ which vanishes exactly on E and such that $f(T) := \sum_{n=0}^{+\infty} \widehat{f}(n)T^n = 0$.

Proof. Let ω be the weight defined by $\omega(n) = \big\|T^n\big\|$ $(n \in \mathbb{Z})$. Let Φ be the continuous morphism from $A_{\omega}(\mathbb{T})$ to $\mathcal{L}(X)$ defined by

$$\Phi(f) = f(T) = \sum_{n=-\infty}^{+\infty} \widehat{f}(n)T^n \qquad \big(f \in A_{\omega}(\mathbb{T})\big).$$

Since the algebra $A_{\omega}(\mathbb{T})$ is regular, we have $\big\{z \in \mathbb{T} :\ f(z) = 0 \quad (f \in \text{Ker } \Phi)\big\} \subset E$ (see [7], theorem 2.5), and so $J_{\omega}(E) \subset \text{Ker } \Phi$. Then the result follows from lemmas 7.1 and 7.2 of [5]. $\square$

Theorem 2.3. *Let E be a K-set, and let s be a nonnegative real number. Then, any invertible operator T on a Banach space with spectrum included in E that satisfies*

$$\begin{aligned} &\big\|T^n\big\| = O\big(n^s\big),\ n \to +\infty\\ \textit{and}\quad &\big\|T^{-n}\big\| = O\big(e^{n^{\beta}}\big),\ n \to +\infty \textit{ for some } \beta < \frac{\delta(E)}{1+\delta(E)}, \end{aligned}$$

also satisfies the stronger property

$$\big\|T^{-n}\big\| = O\big(n^{s+\frac{1}{2}+\varepsilon}\big),\ n \to +\infty,$$

for all $\varepsilon > 0$.

Proof. Let $\varepsilon > 0$ and set $\tilde{s} = s + \frac{1}{2} + \varepsilon$. Without loss of generality, we may assume that $\tilde{s}$ is not an integer. Let t a real number, which is not an integer, and satisfies $s + \frac{1}{2} < t < \tilde{s}$ and $[t] = [\tilde{s}]$. According to lemma 2.1, we can define a continuous morphism Φ from $\lambda_t^{+}(\mathbb{T})$ to $\mathcal{L}(X)$ by

$$\Phi(f) = f(T) = \sum_{n=0}^{+\infty} \widehat{f}(n)T^n \qquad \big(f \in \lambda_t^{+}(\mathbb{T})\big).$$

Let $I = \text{Ker } \Phi$, I is a closed ideal of $\lambda_t^{+}(\mathbb{T})$. We denote by S_I its inner factor, that is the greatest common divisor of all inner factors of the non-zero functions in I (see [8] p.85), and

we set, for $0 \leq k \leq [t]$, $h^k(I) = \big\{ z \in \mathbb{T} :\, f(z) = \ldots = f^{(k)}(z) = 0 \quad (f \in I) \big\}$.
F. A. Shamoyan showed in [12] that

$$I = \Big\{ f \in \lambda_t^+(\mathbb{T}) :\, S(I) |\, S(f) \text{ and } f^{(k)} = 0 \text{ on } h^k(I) \text{ for all } 0 \leq k \leq [t] \Big\},$$

where $S(f)$ denotes the inner factor of f and $S(I) |\, S(f)$ means that $S(f)/S(I)$ is a bounded holomorphic function in $\mathbb{D}$. Since $\beta < \dfrac{\delta(E)}{1+\delta(E)}$, there exists $0 < \delta < \delta(E)$ such that $\beta < \dfrac{\delta}{1+\delta}$. We have, by definition of $\delta(E)$, $\displaystyle\int_0^{2\pi} \frac{1}{d(e^{it}, E)^{\delta}} \mathrm{d}t < +\infty$. So we deduce from lemma 2.2 that there exists an outer function $f \in \mathcal{A}^{\infty}(\mathbb{D})$ which vanishes exactly on E and such that $f \in I$. Therefore, we have $S(I) = 1$ and $h^0(I) \subset E$, so that $N_t^+(E) \cap \lambda_t(\mathbb{T}) \subset I$. Now, as $\Lambda_{\tilde{s}}^+(\mathbb{T}) \subset \lambda_t^+(\mathbb{T})$, we can define a continuous morphism Ψ from $\Lambda_{\tilde{s}}^+(\mathbb{T})$ to $\mathcal{L}(X)$ by $\Psi = \Phi_{|_{\Lambda_{\tilde{s}}^+(\mathbb{T})}}$. Using what precedes, we have

$$N_{\tilde{s}}^+(E) \subset \text{Ker } \Psi.$$

So there exists a continuous morphism $\tilde{\Psi}$ from $\Lambda_{\tilde{s}}^+(\mathbb{T})/N_{\tilde{s}}^+(E)$ into $\mathcal{L}(X)$ such that $\Psi = \tilde{\Psi} \circ \pi_{\tilde{s}}^+$, where $\pi_{\tilde{s}}^+$ is the canonical surjection from $\Lambda_{\tilde{s}}^+(\mathbb{T})$ to $\Lambda_{\tilde{s}}^+(\mathbb{T})/N_{\tilde{s}}^+(E)$. Since E is a K-set, by a theorem of E. M. Dyn'kin [2], it is an interpolating set for $\Lambda_{\tilde{s}}^+(\mathbb{T})$, so that the canonical imbedding i from $\Lambda_{\tilde{s}}^+(\mathbb{T})/N_{\tilde{s}}^+(E)$ into $\Lambda_{\tilde{s}}(\mathbb{T})/N_{\tilde{s}}(E)$ is onto. We have, for $n \geq 0$,

$$T^{-n} = \tilde{\Psi} \circ i^{-1} \circ \pi_{\tilde{s}}(\alpha^{-n}),$$

where $\pi_{\tilde{s}}$ denote the canonical surjection from $\Lambda_{\tilde{s}}(\mathbb{T})$ to $\Lambda_{\tilde{s}}(\mathbb{T})/N_{\tilde{s}}(E)$ and where $\alpha : z \to z$ is the identity map. So we have, for $n \geq 0$,

$$\begin{aligned} \big\| T^{-n} \big\| &\leq \big\| \tilde{\Psi} \circ i^{-1} \big\| \big\| \pi_{\tilde{s}}(\alpha^{-n}) \big\|_{\Lambda_{\tilde{s}}} \\ &\leq \big\| \tilde{\Psi} \circ i^{-1} \big\| (1+n)^{\tilde{s}}, \end{aligned}$$

which completes the proof. □

We give two immediate corollaries of this theorem.

Corollary 2.4. *Let $\xi \in \big(0, \frac{1}{2}\big)$ and let s be a nonnegative real number. Then, any invertible operator T on a Banach space with spectrum included in E_ξ that satisfies*

$$\begin{aligned} &\big\| T^n \big\| = O\big(n^s\big),\ n \to +\infty \\ \textit{and}\quad &\big\| T^{-n} \big\| = O\big(e^{n^{\beta}}\big),\ n \to +\infty \textit{ for some } \beta < b(\xi), \end{aligned}$$

also satisfies the stronger property

$$\big\| T^{-n} \big\| = O\big(n^{s + \frac{1}{2} + \varepsilon}\big),\ n \to +\infty,$$

for all $\varepsilon > 0$.

Proof. It is well known that E_ξ is a K-set (see proposition 2.5 of [5]). Moreover, E_ξ satisfies $\displaystyle\int_0^{2\pi} \frac{1}{d(e^{it},E)^\delta}\mathrm{d}t < +\infty$ if and only if $\delta < 1+\dfrac{\log 2}{\log \xi}$. Indeed, the condition $\displaystyle\int_0^{2\pi} \frac{1}{d(e^{it},E)^\delta}\mathrm{d}t <$ $+\infty$ is equivalent to $\sum\limits_{n=1}^{+\infty}\sum\limits_{i=1}^{2^{n-1}} \left|L_{n,i}\right|^{1-\delta}$, where $L_{n,i}$ are the arcs contiguous to E_ξ, and $\left|L_{n,i}\right|$ are their length, which is equal to $2\pi\xi^{n-1}(1-2\xi)$ (see [10] for further details). Then it is easily seen that the last series converges if and only if $\delta < 1+\dfrac{\log 2}{\log \xi}$, so $\delta(E_\xi) = 1+\dfrac{\log 2}{\log \xi}$. Now, the result follows immediately from theorem 2.3. □

Then we obtain an other immediate result, which generalizes theorem 4.1 of [3]. Indeed, the condition "$\left\|T^{-n}\right\| = O\big(e^{n^\beta}\big)$, $n \to +\infty$" which appears in the following corollary is weaker than the condition used by the authors of [3].

Corollary 2.5. *Let E be a K-set, and let s be a nonnegative real number. Then, there exists a constant $\beta > 0$ independent of s such that any invertible operator T on a Banach space with spectrum included in E that satisfies*

$$\begin{aligned} &\left\|T^n\right\| = O\big(n^s\big),\ n \to +\infty \\ \textit{and}\quad &\left\|T^{-n}\right\| = O\big(e^{n^\beta}\big),\ n \to +\infty, \end{aligned}$$

also satisfies the stronger property

$$\left\|T^{-n}\right\| = O\big(n^{s+\frac{1}{2}+\varepsilon}\big),\ n \to +\infty,$$

for all $\varepsilon > 0$.

Proof. As E is a K-set, we deduce from [2] (section 5, corollary) that $\delta(E) > 0$. Then the result follows immediately from theorem 2.3, with any $\beta < \frac{\delta(E)}{1+\delta(E)}$. □

Remark 2.6:

1) Some results concerning operators with countable spectrum are obtained in [13] and in [1]. Let E be a closed subset of $\mathbb{T}$ and let s, t be two nonnegative reals. We denote by $P(s,t,E)$ the following property: every invertible operator T on a Banach space such that $\mathrm{Sp}\,T \subset E$ and satisfies the conditions:

$$\begin{aligned} &\left\|T^n\right\| = O(n^s)\ (n \to +\infty) \\ &\left\|T^{-n}\right\| = O(e^{\varepsilon\sqrt{n}})\ (n \to +\infty),\ \text{ for all } \varepsilon > 0, \end{aligned}$$

also satisfies the stronger property

$$\left\|T^{-n}\right\| = O(n^t)\,(n \to +\infty).$$

M. Zarrabi showed in [13] (théorème 3.1 and remarque 2.a) that a closed subset E of $\mathbb{T}$ satisfies $P(0,0,E)$ if and only if E is countable. Notice that E is called a Carleson set if $\displaystyle\int_0^{2\pi} \log^+ \frac{1}{d(e^{it},E)}\mathrm{d}t < +\infty$. If E is a countable closed subset of $\mathbb{T}$, we show in [1] that the following conditions are equivalent:

(i) there exist two positive constants C_1, C_2 such that for every arc $I \subset \mathbb{T}$,

$$\frac{1}{|I|}\int_I \log^+ \frac{1}{d(e^{it}, E)}\mathrm{d}t \leq C_1 \log \frac{1}{|I|} + C_2.$$

(ii) E is a Carleson set and for all $s \geq 0$, there exists t such that $P(s,t,E)$ is satisfied. For contractions with spectrum satisfying the Carleson condition, we can see [11].

2) When $\xi = \dfrac{1}{q}$, the constant $b(\dfrac{1}{q})$ in corollary 2.4 is the best possible in view of [6], where the authors built a contraction T such that $\lim\limits_{n\to+\infty} \log \|T^{-n}\| = +\infty$, $Sp\,T \subset E_{\frac{1}{q}}$ and $\log \|T^{-n}\| = O\big(n^{b(\frac{1}{q})}\big)$. According to theorem 6.4 of [4], T doesn't satisfy $\|T^{-n}\| = O\big(n^s\big)$ for any real $s \geq 0$.

References

[1] C. Agrafeuil, *Idéaux fermés de certaines algèbres de Beurling et application aux opérateurs à spectre dénombrable*, preprint.

[2] E. M. Dynkin, *Free interpolation set for Hölder classes*, Mat. Sbornik, **109** (1979), 107-128.

[3] O. El-Fallah et K. Kellay, *Sous-espaces biinvariants pour certains shifts pondérés*, Ann. Inst. Fourier **48** (1998), no. 5, 1543-1558.

[4] J. Esterle, *Uniqueness, strong form of uniqueness and negative powers of contractions*, Banach Center Publ. **30** (1994), 1-19.

[5] J. Esterle, *Distributions on Kronecker sets, strong forms of uniqueness, and closed ideals of A^+*, J. reine angew. Math. **450** (1994), 43-82.

[6] J. Esterle, M. Rajoelina and M. Zarrabi, *On contractions with spectrum contained in Cantor set*, Math. Proc. Camb. Phil. Soc. **117** (1995), 339-343.

[7] J. Esterle, E. Strouse and F. Zouakia, *Theorems of Katznelson-Tzafriri type for contractions*, J. Func. Anal. (2) **94** (1990), 273-287.

[8] K. Hoffman *"Banach spaces of analytic functions"*, Prentic-Hall, Englewood Cliffs, 1962.

[9] J. P. Kahane, *"Séries de Fourier absolument convergentes"*, Erg. Math. **336**, Springer Verlag, Berlin-Heidelberg-New York, 1973.

[10] J. P. Kahane, R. Salem, *"Ensembles parfaits et séries trigonométriques"*, Paris, Hermann, 1963.

[11] K. Kellay, *Contractions et hyperdistributions à spectre de Carleson*, J. London Math. Soc. (2) **58** (1998), 185-196.

[12] F. A. Shamoyan *Closed ideals in algebras of functions analytic in the disc and smooth up to its boundary*, Mat. Sbornik **79** (1994), 425-445.

[13] M. Zarrabi, *Contractions à spectre dénombrable et propriété d'unicité des fermés dénombrables du cercle unité*, Ann. Inst. Fourier **43** (1993), 251-263.

2000 Mathematics Subject Classification: 46J15, 46J20, 47A30.
Key-words: operators, Beurling algebra, spectral synthesis, perfect symmetric set.

AGRAFEUIL Cyril
Cyril.Agrafeuil@math.u-bordeaux.fr
Laboratoire Bordelais d'Analyse et Géométrie (LaBAG), CNRS-UMR 5467 Université Bordeaux I
351, cours de la libération
33405 Talence cedex, FRANCE.